

Prescription Information

This treatment plan contains important information regarding the medication that was prescribed to you by your doctor. The following includes: ingredient information, side effects, dosing instructions, proper storage, and other information important for you to know before starting treatment.

What is the medication used for?

StaminaRx is used to support male sexual performance by helping improve control over ejaculation, prolong the duration of intimacy, and enhance erectile function. The exact dose and whether Sildenafil, Tadalafil, and Apomorphine are included in your treatment will depend on what is the best fit for your symptoms and goals, as determined by your doctor.

Active Prescription Ingredient:

Paroxetine, with or without Sildenafil, Tadalafil, and Apomorphine

What is StaminaRx?

Paroxetine increases serotonin activity in the nervous system. Serotonin plays a key role in controlling ejaculation. By enhancing serotonin signaling, paroxetine helps: delay the ejaculatory reflex, increase intravaginal ejaculatory latency time (IELT), improve control over climax, reduce performance-related anxiety (in some men). Improvement may begin within several days, but optimal results typically develop after 1-3 weeks of consistent use. The optional addition of Sildenafil, Tadalafil, and Apomorphine may help treat erectile dysfunction by relaxing penile muscles and increasing blood flow, promoting stronger, longer-lasting erections. Effects can begin in as little as 15-30 minutes.

What is the best way to use this medication?

Take paroxetine exactly as prescribed by your doctor. Limit to one dose every 24 hours. It can be taken with or without food, but a high-fat meal may delay effects (with Sildenafil, Tadalafil). Avoid alcohol and grapefruit while taking StaminaRx, as they can cause side effects like dizziness or low blood pressure.

What if I take too much StaminaRx?

Taking more StaminaRx than prescribed can increase the risk of serious side effects, including dangerously high or low blood pressure, a prolonged erection lasting over 4 hours (priapism), severe drowsiness, tremors, rapid heartbeat, vomiting, confusion, or seizures. If you've taken too much or experience severe side effects, seek emergency medical attention immediately.

What should I tell my doctor before starting this medication?

Before taking StaminaRx make sure your doctor knows about any of the following health conditions: heart problems (chest pain, a heart rhythm disorder, heart attack), high or low blood pressure, blood circulation problems, retinitis pigmentosa (an inherited condition of the eye), blindness in one or both eyes, bleeding problems, a stomach ulcer, pulmonary veno-occlusive disease (PVOD), liver or kidney disease, a blood disorder, such as sickle cell anemia, multiple myeloma, or leukemia, psychosis or psychotic disorder, Parkinson's disease, a physical deformity of the penis (such as Peyronie's disease), mental health disorders including depression, or if you have been told you should not have sexual intercourse for health reasons. Also, make sure your doctor knows about all of your current medications, supplements, and any other conditions.

What are the possible side effects of this medication?

Common side effects of Paroxetine may include dizziness, headache, nausea, indigestion, anxiety, irritability, agitation, tingling or numbness, changes in sexual function, sweating, flushing, fatigue, vivid dreams, nasal congestion, vision changes, difficulty concentrating, tremors, ringing in the ears, and dry mouth. Sildenafil and Tadalafil may cause headache, indigestion, back or muscle pain, flushing, and nasal congestion. Apomorphine may lead to nausea, dizziness, drowsiness, and discomfort or numbness in the mouth or throat. Serious but rare side effects include: suicidal thoughts (particularly in younger adults), severe mood changes, serotonin syndrome (confusion, rapid heart rate, fever, muscle stiffness), severe allergic reaction. Seek immediate medical attention if serious symptoms occur.

Allergies

Do not take this medication if you are allergic to paroxetine or similar medications in the same class. If your formula includes Sildenafil, Tadalafil, or Apomorphine, avoid use if you are allergic to any of these ingredients or related drugs in their respective classes. Get emergency medical help if you have signs of an allergic reaction: hives, difficulty breathing, swelling of your face, lips, tongue, or throat.

How should I store this medication?

Store this medication in the original container in a cool, dry environment, at a controlled room temperature (68F-77F). Keep out of reach of children and pets.

Pharmacy Disclaimer

The drug and medical information provided by the Pharmacy is not intended to be comprehensive and does not replace the professional judgment of a licensed Healthcare Provider. The Pharmacy is separate and independent from any Healthcare Provider, including telehealth providers, who may prescribe medications dispensed by the Pharmacy. While reasonable efforts are made to ensure information is accurate and current, the Pharmacy makes no guarantees regarding its completeness or timeliness and assumes no responsibility for its use. The absence of a warning for any medication or supplement does not imply safety, effectiveness, or appropriateness for any patient. Questions regarding diagnosis or treatment should be directed to your prescribing Healthcare Provider. Questions regarding dispensing may be directed to the Pharmacy or your pharmacist. Compounded medications are prepared pursuant to a valid prescription and have not been evaluated or approved by the FDA for safety or efficacy. All active ingredients are FDA-approved and compounded in accordance with applicable USP standards.



Prescription Information

STAMINARX



941-921-6645 | 855-610-7227



patientcare@buxhealthcare.com



www.familypharmacy.org

Please scan this code to see our HIPAA Privacy Policy & to find out more detailed information about your product.

