

Prescription Information

This treatment plan contains important information regarding the medication that was prescribed to you by your doctor. The following includes: ingredient information, side effects, dosing instructions, proper storage, and other information important for you to know before starting treatment.

What is the medication used for?

Super ED is designed to treat erectile dysfunction by addressing both the physical and psychological factors that can impact sexual performance. It's innovative 4-in-1 formula blends fast-acting and long-lasting ingredients to help you achieve and sustain firm erections.

By combining the extended effectiveness of tadalafil, the quick response of vardenafil, and the neuro-enhancing effects of apomorphine and oxytocin, this RDT offers a complete, all-in-one solution. It is uniquely formulated to support not just immediate results, but also ongoing sexual well-being—making it a standout option in ED treatment.

Active Prescription Ingredient:

Vardenafil, Tadalafil, Apomorphine, Oxytocin, PT-141 (Bremelanotide), Vitamin D

What is Super ED?

Super ED is a powerful 4-in-1 formula of brain and body-enhancing ingredients, designed to help you achieve and maintain a firm erection, while also delivering stronger and longer-lasting results. Unlike typical ED treatments, this comprehensive RDT combines both fast-acting and extended-release components to support immediate performance and long-term sexual health. Each ingredient targets specific aspects of ED, offering a holistic approach: Tadalafil provides long-lasting support—up to 36 hours—by improving blood flow to the penis, giving you flexibility and confidence without the pressure of perfect timing. Vardenafil works quickly (usually within 30–60 minutes) to enhance blood flow and help you get and keep an erection.

Apomorphine is a dopamine agonist that stimulates receptors in the brain to boost sexual response and blood flow.

Oxytocin, the “love hormone,” may enhance emotional connection, trust, and relaxation—supporting the psychological side of intimacy.

PT-141 (Bremelanotide) is a peptide in the medication class of melanocortin receptors agonists. PT-141 can stimulate neurons in the central nervous system and may help some men with erectile dysfunction see an increase in erectile activity.

Vitamin D - There have been associations between low vitamin D levels and the occurrence or severity of erectile dysfunction. Since many people may be low in vitamin D without knowing it, including a moderate amount in this formulation may help reduce vitamin D deficiencies that could negatively affect erectile performance.

What is the best way to use this medication?

Take this medication by mouth 15-30 minutes before sexual activity. Place under the tongue and dissolve, do not swallow whole. Super ED may be taken with or without food, although when taken with a high-fat meal it may take longer to work. Do not take more than once daily. Avoid drinking alcohol when using this medication. If you have overdosed or have an

erection lasting longer than 4 hours call your doctor and get medical treatment immediately.

What should I tell my doctor before starting this medication?

Tell your doctor if you have ever had: heart problems (chest pain, a heart rhythm disorder, heart attack), high or low blood pressure, blood circulation problems, retinitis pigmentosa (an inherited condition of the eye), blindness in one or both eye, bleeding problems, a stomach ulcer, pulmonary veno-occlusive disease (PVOD), liver or kidney disease, a blood disorder, such as sickle cell anemia, multiple myeloma, or leukemia, psychosis or psychotic disorder, Parkinson's disease, a physical deformity of the penis (such as Peyronie's disease), or if you have been told you should not have sexual intercourse for health reasons. Tell your doctor about all your current medication especially: nitroglycerin, isosorbide dinitrate, isosorbide mononitrate, medicines to treat erectile dysfunction, pulmonary arterial hypertension, antibiotic or antifungal medicine, antiviral medicine to treat hepatitis C or HIV/AIDS, drugs to treat high blood pressure or prostate disorder, seizure medicine, or any medicines that treat nausea or irritable bowel syndrome (IBS).

What are the possible side effects of this medication?

Common side effects may include nausea, dizziness, sleepiness, headache, flushing, nasal congestion, indigestion, upset stomach, and mouth discomfort such as swelling, pain, or sores. Contact your doctor if they persist or become bothersome.

Stop taking Super ED and get emergency medical help if you have: heart attack symptoms; chest pain or pressure, pain spreading to your jaw or shoulder, severe nausea, sweating, vision changes, blurriness, or sudden vision loss, your erection is painful or lasts longer than 4 hours (prolonged erections can damage the penis), ringing in your ears or sudden hearing loss, irregular heartbeat, swelling in your hands, ankles, or feet; shortness of breath, seizure (convulsions), or a light-headed feeling, like you might pass out, high blood pressure- severe headache, pounding in your neck and ears, confusion, severe weakness, unsteady feeling, sleepiness or falling asleep during the day, hallucinations or

psychotic-like behavior, and/or strong (intense) urges.

Allergies

Do not take this medication if you are allergic to Tadalafil or other PDE5 inhibitors, Oxytocin, PT-141, or vitamin D. Get emergency medical help if you have signs of an allergic reaction: hives, difficulty breathing, swelling of your face, lips, tongue, or throat.

How should I store this medication?

Store this medication in its original container in the refrigerator at a temperature between 36°F and 46°F (2°C–8°C). Do not freeze. Keep all medications in a secure location out of reach of children and pets.

Can I use the tablet if it's broken or chipped?

The compound medication is handmade in the pharmacy laboratory and during the shipping process, minor damage may occur. You can still use the tablet if this has occurred.

Pharmacy Disclaimer

The drug and medical information provided by the Pharmacy is not intended to be comprehensive and does not replace the professional judgment of a licensed Healthcare Provider. The Pharmacy is separate and independent from any Healthcare Provider, including telehealth providers, who may prescribe medications dispensed by the Pharmacy. While reasonable efforts are made to ensure information is accurate and current, the Pharmacy makes no guarantees regarding its completeness or timeliness and assumes no responsibility for its use. The absence of a warning for any medication or supplement does not imply safety, effectiveness, or appropriateness for any patient. Questions regarding diagnosis or treatment

should be directed to your prescribing Healthcare Provider. Questions regarding dispensing may be directed to the Pharmacy or your pharmacist.

Compounded medications are prepared pursuant to a valid prescription and have not been evaluated or approved by the FDA for safety or efficacy. All active ingredients are FDA-approved and compounded in accordance with applicable USP standards.



Prescription Information

SUPER ED



941-921-6645 | 855-610-7227



patientcare@buxhealthcare.com



www.familypharmacy.org

Please scan this code to see our HIPAA Privacy Policy & to find out more detailed information about your product.

