

Prescription Information

This treatment plan contains important information regarding the medication that was prescribed to you by your doctor. The following includes: ingredient information, side effects, dosing instructions, proper storage, and other information important for you to know before starting treatment.

What is this medication used for?

Tadalafil is commonly used for erectile dysfunction or increased blood pressure in the lungs. In combination with sexual stimulation tadalafil works by increasing blood flow to the penis to help a man achieve and maintain an erection. Tadalafil is also used to treat the symptoms of an enlarged prostate (benign prostatic hyperplasia-BPH). It helps to relieve symptoms of BPH such as difficulty in beginning the flow of urine, weak stream, and the need to urinate frequently or urgently (including during the middle of the night). Tadalafil

is thought to work by relaxing the smooth muscle in the prostate and bladder.

Active Prescription Ingredient

Tadalafil

Common Brand Names:

Cialis

Adcirca

What is the best way to use this medication?

Take this medication by mouth before sexual activity. Tadalafil may be used up to 36 hours before sexual activity. Tadalafil may be taken with or without food.

Do not break or split tablets.

Do not take more than once daily!

If you take too much or have an erection lasting longer than 4 hours call your doctor and get medical treatment immediately.

What should I tell my doctor before starting this medication?

Tell your doctor if you have ever had: heart problems (chest pain, a heart rhythm disorder, heart attack), high or low blood pressure, blood circulation problems, retinitis pigmentosa (an inherited condition of the eye), one or both eyes, bleeding blindness in problems, a stomach ulcer, pulmonary veno-occlusive disease (PVOD), liver or kidney disease, a blood cell disorder such as sickle cell anemia, multiple myeloma or leukemia, a physical deformity of the penis (such as Peyronie's disease), or if you have been told you should not have sexual intercourse for health reasons. Tell your doctor about all your medicines, especially: nitroglycerin, current isosorbide dinitrate, isosorbide mononitrate, treat erectile dysfunction or medicines to pulmonary arterial hypertension, antibiotic or antifungal medicine, antiviral medicine to treat hepatitis C or HIV/ AIDS, drugs to treat high blood pressure or a prostate disorder, or seizure medicine. Nitrates are also found in some recreational drugs such as amyl nitrate or nitrite ("poppers").

What are the possible side effects of this medication?

Stop taking tadalafil and get emergency medical help if you have: heart attack symptoms - chest pain or pressure, pain spreading to your jaw or shoulder, nausea, sweating, vision changes or sudden vision loss; an erection that is painful or lasts longer than 4 hours (prolonged erection can damage the penis); ringing in your ears or sudden hearing loss; irregular heartbeat; swelling in your hands, ankles, or feet; shortness of breath, seizure (convulsions), or a light-headed feeling like you might pass out.

Common Tadalafil side effects may include: flushing (warmth, redness, or tingly feeling), headache, dizziness, heartburn, nausea or upset stomach, abnormal vision (blurred vision, changes in color vision), runny or stuffy nose, nosebleeds, sleep problems (insomnia), muscle pain, or back pain.

Allergies

Do not take this medication if you are allergic to tadalafil or other PDE5 inhibitors. Get emergency medical help if you have signs of an allergic reaction: hives, difficulty breathing, swelling of your face, lips, tongue, or throat.

How should I store this medication?

Store this medication in the original container in a cool, dry environment, at a controlled room temperature (68F-77F).

FAQ's

Please visit the Family Pharmacy website FAQ section or search the <https://www.familypharmacy.org/our-blog-2> for more informative articles and literature.

Pharmacy Disclaimer

The drug and medical information provided by the Pharmacy is not intended to be comprehensive and does not replace the professional judgment of a licensed Healthcare Provider. The Pharmacy is separate and independent from any Healthcare Provider, including telehealth providers, who may prescribe medications dispensed by the Pharmacy.

While reasonable efforts are made to ensure information is accurate and current, the Pharmacy makes no guarantees regarding its completeness or timeliness and assumes no responsibility for its use. The absence of a warning for any medication or supplement does not imply safety, effectiveness, or appropriateness for any patient.

Questions regarding diagnosis or treatment should be directed to your prescribing Healthcare Provider. Questions regarding dispensing may be directed to the Pharmacy or your pharmacist.

Compounded medications are prepared pursuant to a valid prescription and have not been evaluated or approved by the FDA for safety or efficacy. All active ingredients are FDA-approved and compounded in accordance with applicable USP standards.



Prescription Information

TADALAFIL



941-921-6645 | 855-610-7227



patientcare@buxhealthcare.com



www.familypharmacy.org

Please scan this code to see our HIPAA Privacy Policy & to find out more detailed information about your product.

